


Ch07

Carboxylic Acids


Combining the hydroxyl and carbonyl functional groups.
To make organic acids.



Important Dates



- ▶ This Wednesday:
 - Lab Checkout (you must check out of your lab locker or you may be fined)
 - Lab Practical, last lab quiz (25 points)
 - Last day to turn in past due labs and homework (start of lab period)
- ▶ Mon, March 21st: Final Exam **9:15-11:15 a.m.**
you **must** take the final exam to pass the course
Final Exam is worth 120 points
Final Exam is cumulative
- ▶ Mon, April 4th: Final Grades will be submitted to the college.
(They won't be available until they are posted;
they will be posted as soon as they are available.)



Carboxylic Acids & Esters



Carboxylic Acids

- ▶ Carboxyl Group
 - ▶ Compound Functional Groups
- ▶ Properties & Structure
 - ▶ IM Forces
 - ▶ Acid-Base
- ▶ Naming
 - ▶ Carboxylic Acids
 - ▶ Carboxylic Acid Salts



Mixing Functional Groups

- ▶ Adding/Changing Properties
- ▶ Naming Substances w/ multiple functional groups
- ▶ Willow Bark

Esters

- ▶ Structure & Properties
- ▶ Naming

Reactions

- ▶ Esterification
- ▶ Hydrolysis
- ▶ Saponification




Carboxylic Acids

- ▶ Carboxylic Acids have many interesting properties.
 - ▶ Many of the sharp flavors we enjoy in food (vinegar, grapefruit, lemon...) are produced by simple substances of the carboxylic acid family of organic substances.
 - ▶ These organic molecules are acids. Like the simple binary acids you're already familiar with (HCl, HBr) they release free protons (H^+).
 - ▶ These substances have high acidity (low pH).
 - ▶ They neutralize bases like NaOH to form water.
 - ▶ Many toxins are simple carboxylic acids, like the formic acid in ant venom.
 - ▶ Salts formed from carboxylic acids cause food to be resistant to mold and microorganisms that cause spoilage.
 - ▶ They play important roles in how animals store and process energy.




Carboxyl Functional Group

- ▶ Carboxylic Acids are defined by the **carboxyl functional group**.
 - ▶ The carboxyl group is a composite of two functional groups you're already familiar with.
 - ▶ The hydroxyl group.
 - ▶ The carbonyl group.





Carbonyl group







Hydroxyl group

Carboxyl group



Carboxyl Functional Group

- ▶ Carboxylic Acids are defined by the **carboxyl functional group**.
 - ▶ The carboxyl group is a composite of two functional groups you're already familiar with.
 - ▶ The hydroxyl group.
 - ▶ The carbonyl group.
 - ▶ Both of those groups are polar.
 - ▶ Carboxylic acids are strongly polar because they have two polar groups.



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification




▶ Hydrolysis

▶ Saponification



Carboxyl Functional Group



- ▶ Carboxylic Acids are defined by the **carboxyl functional group**.
 - ▶ The carboxyl group is a composite of two functional groups you're already familiar with.
 - ▶ The hydroxyl group.
 - ▶ The carbonyl group.
 - ▶ Both of those groups are polar.
 - ▶ Carboxylic acids are strongly polar because they have two polar groups.
 - ▶ Each carboxylic acid forms hydrogen bonds with many water molecules at once.
 - ▶ Carboxylic acids with up to four carbon atoms are very soluble in water.
 - ▶ As the number of carbons increases, the solubility of the carboxylic acid in water is reduced.




Properties & Structure

- ▶ Carboxylic Acids are defined by the **carboxyl functional group**.
 - ▶ The carboxyl group is a composite of two functional groups you're already familiar with.
 - ▶ The hydroxyl group.
 - ▶ The carbonyl group.
 - ▶ Both of those groups are polar.
 - ▶ Carboxylic acids are strongly polar because they have two polar groups.
 - ▶ Each carboxylic acid forms hydrogen bonds with many water molecules at once.
 - ▶ Carboxylic acids with up to four carbon atoms are very soluble in water.
 - ▶ As the number of carbons increases, the solubility of the carboxylic acid in water is reduced.

| IUPAC Name | Condensed Structural Formula | Solubility in Water |
|----------------|---|---------------------|
| Methanoic acid | $\text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Soluble |
| Ethanoic acid | $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Soluble |
| Propanoic acid | $\text{CH}_3-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Soluble |
| Butanoic acid | $\text{CH}_3-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Soluble |
| Pentanoic acid | $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Soluble |
| Hexanoic acid | $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Slightly soluble |
| Benzoic acid | $\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Slightly soluble |



Properties & Structure



▶ Carboxylic Acids experience all three kinds of intermolecular forces:

- ▶ Van der Waals Forces
- ▶ Dipole-Dipole Forces
- ▶ Hydrogen Bonding

- ▶ Because carboxylic acids have both a hydroxyl and carbonyl group they form more hydrogen bonds than alcohols.
- ▶ Carboxylic Acids have higher boiling points and melting points than even alcohols.



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification




▶ Hydrolysis

▶ Saponification




Properties & Structure

- ▶ Carboxylic Acids are defined by the **carboxyl functional group**.
 - ▶ Carboxylic acids are strongly polar because they have two polar groups.
 - ▶ The two polar groups cooperate to balance negative charge, stabilizing the molecule enough to allow it to release protons.
 - ▶ To act as an organic acid.




H^+




Properties & Structure

- ▶ Carboxylic Acids are defined by the **carboxyl functional group**.
 - ▶ Carboxylic acids are strongly polar because they have two polar groups.
 - ▶ The two polar groups cooperate to balance negative charge, stabilizing the molecule enough to allow it to release protons.
 - ▶ To act as an organic acid.



| Functional group | Example | pKa | Conjugate Base |
|---------------------|--|-------|--|
| Alkane | $\text{H}_3\text{C}-\text{CH}_2-\text{CH}_3$ | ~50 | $\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2^-$ |
| Alkene | $\text{H}_2\text{C}=\text{CH}_2$ | ~43 | $\text{H}_2\text{C}=\text{CH}^-$ |
| Ketone/ aldehyde | $\text{H}_3\text{C}-\text{C}(=\text{O})-\text{CH}_3$ | 20-24 | $\text{H}_3\text{C}-\text{C}(=\text{O})-\text{CH}_2^-$ |
| Alcohol | $\text{H}_3\text{C}-\text{OH}$ | 17 | $\text{H}_3\text{C}-\text{O}^-$ |
| Water | $\text{HO}-\text{H}$ | 16 | HO^- |
| Thiols | $\text{CH}_3\text{S}-\text{H}$ | 13 | CH_3S^- |
| Carboxylic acids | $\text{H}_3\text{C}-\text{C}(=\text{O})-\text{OH}$ | 4 | $\text{H}_3\text{C}-\text{C}(=\text{O})-\text{O}^-$ |
| Sulfuric acid | H_2SO_4 | -3 | HSO_4^- |
| Hydrochloric acid | HCl | -6 | Cl^- |



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base



Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification





▶ Hydrolysis


▶ Saponification



Naming Carboxylic Acids


- ▶ The simplest carboxylic acids are formic acid and acetic acid.
- ▶ Carboxylic acids are named with IUPAC using the family suffix **-oic acid**.

| Condensed Structural Formula | IUPAC Name | Common Name | Ball-and-Stick Model |
|---|----------------|----------------|---|
| $\text{H}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Methanoic acid | Formic acid |  |
| $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Ethanoic acid | Acetic acid |  |
| $\text{CH}_3-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Propanoic acid | Propionic acid |  |
| $\text{CH}_3-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{OH}$ | Butanoic acid | Butyric acid |  |



Naming Carboxylic Acids


- ▶ The simplest carboxylic acids are formic acid and acetic acid.
- ▶ Carboxylic acids are named with IUPAC using the family suffix **-oic acid**.




3-Methylpentanoic acid



3-Hexynoic acid





trans-3-Chloro-5-heptenoic acid




Carboxylate Ions

- ▶ Carboxylic Acids are defined by the **carboxyl functional group**.
 - ▶ Carboxylic acids are strongly polar because they have two polar groups.
 - ▶ The two polar groups cooperate to balance negative charge, stabilizing the molecule enough to allow it to release protons.
 - ▶ To act as an organic acid.




Carboxylic Acid




Ethanoic acid
(acetic acid)

Carboxylate Ion



Ethanoate ion
(acetate ion)


Hydronium
ion





Carboxylate Ions

- ▶ The ions formed from carboxylic acids are named the same way.
- ▶ We use the family suffix **-oate ion** to name the carboxylate ion formed.

Carboxylic Acid




Carboxylate Ion





Carboxylate Ions

- ▶ The ions formed from carboxylic acids are named the same way.
- ▶ We use the family suffix **-oate ion** to name the carboxylate ion formed.
- ▶ You already know how to name the salts formed from ions (organic or otherwise).
- ▶ Just combine the name of the two ions.


Carboxylic Acid




Carboxylate Ion




Examples



Potassium *trans*-2,3-Dichloro-4-hexenoate







Lithium 4-Bromo-3-butynoate



Examples

▶ Carboxylic acid salts are used as preservatives and flavor enhancers such as...

- * sodium propanoate, which is used in cheese and breads
- * sodium benzoate, which inhibits growth of mold and bacteria and is added to fruit juices, margarine, relishes, salads, and jams
- * monosodium glutamate, MSG, which is added to meats, fish, vegetables, and baked goods to enhance flavor



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts



Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification


▶ Hydrolysis

▶ Saponification




Mixing Functional Groups


- ▶ As structures get larger, they will start containing more than one functional group.
 - ▶ Carboxylic acids contain both **carbonyl** and **hydroxyl** groups.
 - ▶ When they're connected we often describe the combination as a new functional group, the **carboxyl** group in this case.
- ▶ The functional groups will interact, creating new dynamics within the molecule and therefore new properties.
 - ▶ The **hydroxyl** group behaves *differently* in the alcohol, phenol, and carboxylic families.




Carbonyl group



Carboxyl group




Hydroxyl group




Mixing Functional Groups

- ▶ As structures get larger, they will start containing more than one functional group.
 - ▶ Carboxylic acids contain both **carbonyl** and **hydroxyl groups**.
 - ▶ When they're connected we often describe the combination as a new functional group, the **carboxyl group** in this case.
- ▶ The functional groups will interact, creating new dynamics within the molecule and therefore new properties.
 - ▶ The **hydroxyl group** behaves *differently* in the alcohol, phenol, and carboxylic families.



Alcohol Family



Phenol Family




Carboxylic Acid Family



Mixing Functional Groups

- ▶ As structures get larger, they will start containing more than one functional group.
 - ▶ Carboxylic acids contain both **carbonyl** and **hydroxyl groups**.
 - ▶ When they're connected we often describe the combination as a new functional group, the **carboxyl group** in this case.
- ▶ The functional groups will interact, creating new dynamics within the molecule and therefore new properties.
 - ▶ The **hydroxyl group** behaves *differently* in the alcohol, phenol, and carboxylic families.


Alcohol Family




pKa = 14-16
(not acidic)

- Somewhat soluble in water.
- High pH (not acidic)



Alcohol Family



Phenol Family




Carboxylic Acid Family




Mixing Functional Groups

- ▶ The functional groups will interact, creating new dynamics within the molecule and therefore new properties.
 - ▶ The **hydroxyl group** behaves *differently* in the alcohol, phenol, and carboxylic families.
 - ▶ Structural features, like unsaturations, can change it's behavior and therefore the properties of the substance.



Phenol Family



$pK_a = 10$
(slightly acidic)


- More soluble in water.
- Moderate pH.



Mixing Functional Groups



- ▶ The functional groups will interact, creating new dynamics within the molecule and therefore new properties.
 - ▶ The **hydroxyl group** behaves *differently* in the alcohol, phenol, and carboxylic families.
 - ▶ Structural features, like unsaturations, can change its behavior and therefore the properties of the substance.
 - ▶ Other functional groups, like **carbonyls**, can also interact and change behavior and properties.

Carboxylic Acid Family



pKa = 4
(an acid)


- Very soluble in water.
- High pH.



Mixing Functional Groups

- ▶ The functional groups will interact, creating new dynamics within the molecule and therefore new properties.
 - ▶ The **hydroxyl group** behaves *differently* in the alcohol, phenol, and carboxylic families.
 - ▶ Structural features, like unsaturations, can change it's behavior and therefore the properties of the substance.
 - ▶ Other functional groups, like **carbonyls**, can also interact and change behavior and properties.


Alcohol Family



pKa = 14-16
(not acidic)

- Somewhat soluble in water.
- High pH (not acidic)


Phenol Family



pKa = 10
(slightly acidic)


- More soluble in water.
- Moderate pH.

Carboxylic Acid Family



pKa = 4
(an acid)

- Very soluble in water.
- Low pH (acidic)



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties



▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification

▶ Hydrolysis


▶ Saponification




Mixing Functional Groups

- ▶ As structures get larger, they will start containing more than one functional group.
 - ▶ Carboxylic acids contain both **carbonyl** and **hydroxyl** groups.
 - ▶ When they're connected we often describe the combination as a new functional group, the **carboxyl** group in this case.
- ▶ When the functional groups are not connected, we need a way to assign a family and address the functional groups that don't define that family.
- ▶ For simple mixed functional group compounds, we assign them to the family of the most oxidized functional group in the molecule.
 - ▶ Aldehydes trump ketones (they can be further oxidized, ketones can't)


How do we assign addresses to the other functional groups?




Name it as a ketone.



Name it as an aldehyde.



Name it as a carboxylic acid.




Mixing Functional Groups


- ▶ For simple mixed functional group compounds, we assign them to the family of the most oxidized functional group in the molecule.
 - ▶ Aldehydes trump ketones (they can be further oxidized, ketones can't)
- ▶ The other functional groups get demoted to substituents and addressed with the same system we've been using all along.

| Functional Group | | Prefix (used as substituent) |
|------------------|----------------|---------------------------------|
| –OH | Hydroxyl Group | hydroxy- |
| –SH | Thiol Group | sulfanyl- |
| –CO– | Carbonyl Group | oxo- |


| Family | Suffix (for family) | Functional Group |
|-----------|------------------------|------------------|
| Alcohols | -ol | –OH |
| Phenols | -phenol | –OH |
| Thiols | -thiol | –SH |
| Ketones | -one | –CO– |
| Aldehydes | -al | –CO– |




Name it as a ketone.




Name it as an aldehyde.




Name it as a carboxylic acid.



Examples




3-Hydroxy-2-pentanone




4-Oxopentanal


| Functional Group | | Prefix (used as substituent) |
|------------------|----------------|---------------------------------|
| -OH | Hydroxyl Group | hydroxy- |
| -SH | Thiol Group | sulfanyl- |
| -CO- | Carbonyl Group | oxo- |



3-Hydroxy-4-oxopentanal



3-Hydroxy-4-oxopentanoic acid



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

→ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification


▶ Hydrolysis

▶ Saponification




Willow Bark


salicylic acid




- ▶ By the 1800s, chemists discovered that salicin from the willow tree bark and leaves was responsible for pain relief, fever reduction and reduced inflammation.
- ▶ The body converts salicin to salicylic acid by reducing the ether and then oxidizing the primary alcohol to a carboxylic acid.
- ▶ **Salicylic acid** is the substance that has analgesic, anti-inflammatory, and antipyretic properties.



[RED]



[OX]




salicylic acid

analgesic,
anti-inflammatory,
and antipyretic





Willow Bark

salicylic acid




- ▶ Salicylic acid is the substance that has analgesic, anti-inflammatory, and antipyretic properties.
- ▶ Salicylic acid is very acidic and can cause upset stomachs, that side effect makes it difficult to use in many cases.
- ▶ Phenols and carboxylic acids are acidic, but other substances with these functional groups don't have the same side effect.




Willow Bark


salicylic acid




- ▶ Salicylic acid is the substance that has analgesic, anti-inflammatory, and antipyretic properties.
- ▶ Salicylic acid is very acidic and can cause upset stomachs, that side effect makes it difficult to use in many cases.
- ▶ Phenols and carboxylic acids are acidic, but other substances with these functional groups don't have the same side effect.
- ▶ In salicylic acid, the functional groups interact to create that greater acidity.




Willow Bark



- ▶ Chemists tried to improve on salicylic acid to reduce that side effect while retaining its other useful properties.
- ▶ The first thing they tried was replacing the hydrogen in the carboxylic acid with a methyl group.
- ▶ Carboxylic Acids and alcohols can react in acid and water to combine. This is a reversible reaction.
- ▶ This is a **condensation** or **esterification** reaction.



salicylic acid




methanol

methyl salicylate



Oil of Wintergreen


methyl salicylate




- ▶ Methyl salicylate (oil of wintergreen) retained many of the properties of salicylic acid.
 - ▶ It's still an analgesic (pain reliever).
- ▶ It lost other properties.
 - ▶ Without the carboxylic acid group it is less acidic.
 - ▶ It's also less water soluble, more greasy.
 - ▶ It's not easily consumed.
- ▶ It has other properties all it's own.
 - ▶ In small quantities it's used as food flavoring, it's responsible for the mint taste in spearmint gum.
 - ▶ It can be absorbed through the skin. It's the active ingredient in Tiger Balm, Icy Hot, and Bengay ointments.
 - ▶ It has a high toxicity. One spoonful is three times a lethal dose.
 - ▶ In 2007 a professional runner died from using too much Bengay.




Willow Bark



- ▶ Chemists tried to improve on salicylic acid to reduce that side effect while retaining its other useful properties.
- ▶ The next thing they tried was replacing the hydrogen in the phenol with an acetal group.
- ▶ Carboxylic Acids and alcohols can react in acid and water to combine. This is a reversible reaction.
- ▶ They used the same esterification reaction to disrupt the phenols contribution to salicylic acids high acidity.



Aspirin



- ▶ Acetylsalicylic acid (aspirin) retained many of the properties of salicylic acid.
 - ▶ It's still an analgesic (pain reliever)
 - ▶ It's also an antipyretic (fever reducer)
 - ▶ It's an anti-inflammatory agent
- ▶ It lost other properties.
 - ▶ It doesn't have the same disruptive effect on the stomach.
- ▶ Bayer chemicals introduced aspirin as a product in 1897 and has been selling it ever since.



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark



Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification


▶ Hydrolysis

▶ Saponification




Esters


- ▶ Both oil of wintergreen and aspirin are examples of another family of organic compounds.
- ▶ **Esters** are substances derived from an acid (organic or inorganic) in which at least one -OH (hydroxyl) group is replaced by an -O-alkyl or -O-aryl group.
- ▶ Usually, esters are derived from a carboxylic acid and an alcohol.
- ▶ Esters comprise most naturally occurring fats and oils.




Aspirin
An Ester



Oil of Wintergreen
An Ester




Salicylic acid
A Carboxylic Acid



Esters

- ▶ Many of the fragrances of perfumes and flowers and the flavors of fruits are due to esters.
- ▶ Simple esters are volatile, so we can smell them, and they are soluble in water, so we can taste them.
 - ▶ Being a hydrogen bond accepter esters are reasonably water soluble.
 - ▶ Having no hydroxy group, they cannot be a hydrogen bond donator, so they have a lower boiling point (more volatile).






| Condensed Structural Formula and Name | Flavor/Odor |
|---|-------------|
| $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_2-\text{CH}_2-\text{CH}_3$ Propyl ethanoate (propyl acetate) | Pears |
| $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3$ Pentyl ethanoate (pentyl acetate) | Bananas |
| $\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3$ Octyl ethanoate (octyl acetate) | Oranges |
| $\text{CH}_3-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_2-\text{CH}_3$ Ethyl butanoate (ethyl butyrate) | Pineapples |
| $\text{CH}_3-\text{CH}_2-\text{CH}_2-\overset{\text{O}}{\parallel}{\text{C}}-\text{O}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3$ Pentyl butanoate (pentyl butyrate) | Apricots |



Esters

- ▶ Esters are used in many ways.
 - ▶ Fragrant esters are used in perfumes, essential oils, food flavorings, and cosmetics.
 - ▶ Natural esters are found in pheromones
 - ▶ Nitrated esters, such as nitroglycerin, are known for their explosive properties.
 - ▶ Polymers formed from esters are called polyesters are used to make plastics and synthetic cloth.
 - ▶ Esters are used to make surfactants (soaps, detergents, cleansers...)



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification

▶ Hydrolysis


▶ Saponification



Esters


- ▶ Simple esters are named with the family suffix **-oate**.
- ▶ The name has two parts.
 - ▶ Start with the underlying carboxylic acid and replace the **-oic acid** with **-oate**. That's the second part.
 - ▶ In front of that put the substituent bonded to the oxygen, as if it were a chain branch.

Carboxylic Acid




Ethanoic acid
(acetic acid)


Ester



Methyl ethanoate
(methyl acetate)



Ethyl butanoate




Propyl ethanoate




Esters

- ▶ Simple esters are named with the family suffix **-oate**.
- ▶ The name has two parts.
 - ▶ Start with the underlying carboxylic acid and replace the **-oic acid** with **-oate**. That's the second part.
 - ▶ In front of that put the substituent bonded to the oxygen, as if it were a chain branch.



Cyclohexyl 3-chloropentanoate



cis-2-Butenyl *trans*-3-pentenoate



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

Reactions

▶ Esterification


▶ Hydrolysis

▶ Saponification





Reactions of Esters

- ▶ Esters are formed by condensing an alcohol and a carboxylic acid.
- ▶ The reaction is also called an **esterification** of a carboxylic acid.
- ▶ It's an equilibrium reaction, but driven forward by using a large excess of the alcohol.





Reactions of Esters

- ▶ **Hydrolysis** of esters is breaking them into alcohols and carboxylic acids.
- ▶ It can be accomplished with acid and heat.
 - ▶ It's an equilibrium reaction, but driven forward by using a large excess water.
 - ▶ It's the reverse of esterification.




Reactions of Esters

- ▶ **Hydrolysis** of esters is breaking them into alcohols and carboxylic acids.
- ▶ Hydrolysis can also be accomplished with strong base.
- ▶ This type of hydrolysis is called saponification (soap making).
- ▶ It produces carboxylic acid salts instead of acids.




Reaction Summary



▶ Esterification:



▶ Hydrolysis:



▶ Saponification:



Carboxylic Acids & Esters

▶ Carboxylic Acids

▶ Carboxyl Group

▶ Compound Functional Groups

▶ Properties & Structure

▶ IM Forces

▶ Acid-Base

▶ Naming

▶ Carboxylic Acids

▶ Carboxylic Acid Salts

▶ Mixing Functional Groups

▶ Adding/Changing Properties

▶ Naming Substances w/ multiple functional groups

▶ Willow Bark

▶ Esters

▶ Structure & Properties

▶ Naming

▶ Reactions

▶ Esterification

▶ Hydrolysis

▶ Saponification



Questions?

